## INFLUENZA WEEKLY UPDATE

2006/39: 25 - 29 September 2006

## **Summary**

Influenza in the population is estimated using the number of GP consultations for flu like illnesses. For 2006 this started on May 1. Current levels are low and the influenza activity this week has decreased compared with the previous week.

The health districts with the highest rate last week were Eastern Bay of Plenty and Otago. 60% of the cases this week were in adults aged between 20 and 64 years and 16% were in children aged 5-19 years.

So far this year 99% of the laboratory tested cases were influenza A with a small number of influenza B cases.

## Thresholds used to describe influenza-like activity\*


Term used		Consultation rate (per 100 000 population)
Baseline		<= 49
Normal	low	50-99
seasonal	moderate	100-149
activity	high	150-249
higher than expected		250-399
severe epidemic		>= 400


<sup>\*</sup>Note: This was published in New Zealand Public Health Report 2001, 8(1):9-12 "Influenza surveillance and immunisation in New Zealand, 1990-1999"

In the past week, a total of 37 consultations for influenza-like illness were reported from 78 general practices in 21 out of 24 health districts. This gives a weekly consultation rate of 11.1 per 100 000 patient population.

Figure 1 compares the consultation rates for influenza-like illness for each health district over the past week. Eastern Bay of Plenty had the highest consultation rate (243.8 per 100 000), followed by Otago (39.6 per 100 000).

Figure 1





[] Health district not participating


Eleven swabs were sent from the sentinel surveillance in the past week. Twelve swabs were received by virology laboratories. Of these, two A/New Caledonia/20/1999 (H1N1)-like were isolated from Wairarapa (1) and Southland (1).

In addition, four influenza A viruses (yet to be sub-typed) were reported this week from the laboratory-based (non-sentinel) surveillance from Otago (2), North West Auckland (1), and Canterbury (1).

Figure 2 shows the cumulative total of influenza isolates confirmed (sentinel and laboratory-based surveillance) to the end of week 39, 29 September 2006. A total of 764 influenza viruses were identified, 381 as A/New York/55/2004 (H3N2) - like, 326 as influenza A (yet to be sub-typed), 52 as A/New Caledonia/20/1999 (H1N1)-like, two as B/Shanghai/361/2002-like, two as B/Malaysia/2506/2004-like, and one as influenza B (yet to be typed).

<sup>\*</sup> Health districts did not participate for the week

Figure 2






Figure 3 shows the weekly national consultation rates for 2004 and 2005 seasons, and 2006 so far.

Figure 3





Figure 4 illustrates consultation rates for influenza-like illness mapped by health district for week 39, 2006.

Figure 4



Compiled by: Liza Lopez

Population and Environmental Health ESR Kenepuru Science Centre

P O Box 50 348 PORIRUA phone: 04 914-0647 fax: 04 914-0770

email: liza.lopez@esr.cri.nz