

**Gonorrhoea : Table 5 - Site of Diagnosis  
January 2003 to March 2003**

|                         | Confirmed                |              |                         |             |                                    |                         |          |                          |          |             |           |              |            |
|-------------------------|--------------------------|--------------|-------------------------|-------------|------------------------------------|-------------------------|----------|--------------------------|----------|-------------|-----------|--------------|------------|
|                         | Uncomplicated infection  |              |                         |             | PID/<br>epididymitis<br>No.¹    %² | Extra-genital infection |          |                          |          | Probable    |           | Total³       |            |
|                         | Urogenital<br>No.¹    %² |              | Anorectal<br>No.¹    %² |             |                                    | Pharynx<br>No.¹    %²   |          | Other site<br>No.¹    %² |          | No.¹    %²  |           |              |            |
| STD Clinics             | 140                      | 81.9%        | 5                       | 2.9%        | 4                                  | 2.3%                    | 5        | 2.9%                     | 2        | 1.2%        | 17        | 9.9%         | 171        |
| Family Planning Clinics | 51                       | 81.0%        | 0                       | 0.0%        | 2                                  | 3.2%                    | 0        | 0.0%                     | 0        | 0.0%        | 10        | 15.9%        | 63         |
| Student Health Clinics  | 3                        | 100.0%       | 0                       | 0.0%        | 0                                  | 0.0%                    | 0        | 0.0%                     | 0        | 0.0%        | 0         | 0.0%         | 3          |
| <b>Total</b>            | <b>194</b>               | <b>82.0%</b> | <b>5</b>                | <b>2.0%</b> | <b>6</b>                           | <b>3.0%</b>             | <b>5</b> | <b>2.0%</b>              | <b>2</b> | <b>1.0%</b> | <b>27</b> | <b>11.0%</b> | <b>237</b> |

<sup>1</sup> No. = no. patients with that diagnosis

<sup>2</sup> % = no. patients with that diagnosis / total no. patients with confirmed or probable Gonorrhoea

<sup>3</sup> Total = total no. patients with confirmed or probable Gonorrhoea; note that patients may have confirmed Gonorrhoea at more than one site