ISSN 1176-7316

Laboratory Surveillance of Chlamydia and Gonorrhoea in New Zealand

October to December 2008

Prepared as part of a Ministry of Health Contract for scientific services

by

STI Surveillance Team Population and Environmental Health Group Institute of Environmental Science & Research Ltd.

February 2009

Client Report FW 09010

Disclaimer

This report or document ("the Report") is given by the Institute of Environmental Science and Research Limited ("ESR") solely for the benefit of the Ministry of Health, Public Health Service Providers and other Third Party Beneficiaries as defined in the Contract between ESR and the Ministry of Health, and is strictly subject to the conditions laid out in that Contract.

Neither ESR nor any of its employees makes any warranty, express or implied, or assumes any legal liability or responsibility for use of the Report or its contents by any other person or organisation.

Acknowledgements

The authors wish to acknowledge that this report could not have been generated without the continuing support of the participating laboratories (see Appendix A).

Members of the STI Surveillance Team at ESR who have contributed to the collection, collation, analysis and reporting of data include Dinusha Bandara, Jonathan Williman, Kerry Sexton and Pauline Quinn.

This report is available on the Internet at <u>www.surv.esr.cri.nz</u> Ad Hoc STI data requests may be emailed to <u>survqueries@esr.cri.nz</u>

Contents

Introduction	1
Methods	2
Chlamydia Results – October to December, 2008	3
Chlamydia Incidence	3
Data tables: Chlamydia Incidence, October to December, 2008 Chlamydia trends: Auckland, Waikato and BOP Regions	4 5
Gonorrhoea Results – October to December, 2008	6
Gonorrhoea Incidence	6
Data tables: Gonorrhoea Incidence, October to December, 2008	7
Gonorrhoea trends: Auckland, Waikato and BOP Regions	8
Appendix A	9

Introduction

In New Zealand, sexually transmitted infections (STIs) are not notifiable and so surveillance efforts rely upon laboratories and clinics voluntarily providing data. Laboratory surveillance of gonorrhoea began in the Auckland, Waikato and Bay of Plenty (BOP) regions in 1998. Laboratory surveillance of chlamydia began in the Waikato and BOP regions in 1998 and in the Auckland region in 2001. Since June 2004, efforts have been made to extend STI surveillance to additional laboratories across New Zealand.

This report presents data on chlamydia and gonorrhoea cases reported in the fourth quarter (October to December) 2008 from thirty-nine laboratories across New Zealand.

Quarterly reports of clinic-based surveillance are provided separately and are available at <u>www.surv.esr.cri.nz</u>.

All results and analyses are based on data submitted prior to the 5th February 2009. Any data submitted after this date is not included in this report due to time constraints. All results should be regarded as provisional.

Methods

The participating laboratories report anonymised data on laboratory-confirmed cases of chlamydia and gonorrhoea, by age and sex. Laboratories only report specimens received directly from health care settings and do not report data on specimens which were subcontracted to their laboratory from outside of their region. The diagnostic tests used by each laboratory differ.

An attempt has been made to de-duplicate the data where one patient may have multiple positive specimens. If this is not possible then it was assumed that each test-positive specimen was equivalent to one test-positive patient. As it is possible for one patient to have more than one positive specimen taken for the one STI episode, the true incidence may be less than that reported here.

STI Rates

Population-based disease rates for the quarter were calculated by dividing the total number of test-positive reported cases for chlamydia and gonorrhoea by the total 'usually resident' population data for the District Health Boards (DHBs) included in the Auckland, Waikato and BOP regions (see below). Population data was supplied from the 2006 Census, Statistics New Zealand.

For several reasons population-based rates have not been determined for the new laboratories. Data submitted from newly participating laboratories is presented as the number of test-positive reported cases by age and sex. As the majority of laboratories within the new DHBs are not, at present, routinely reporting STI data population rates cannot be calculated accurately.

In contrast to reports prepared prior to 2004 STI rates are not annualised.

Auckland region	Includes Auckland DHB and Counties Manukau DHB for calculation of gonorrhoea rates. Includes Auckland DHB, Counties Manukau DHB and Waitemata DHB for calculation of chlamydia rates.
Waikato region	Includes Waikato DHB.
BOP region	Includes BOP DHB and Lakes DHB.

Data completeness

<u>Chlamydia data</u>

Thirty-six laboratories provide chlamydia data for STI surveillance. All 36 laboratories submitted chlamydia data for the fourth quarter (October to December) 2008.

<u>Gonorrhoea data</u>

Thirty-one laboratories provide gonorrhoea data for STI surveillance. All 31 laboratories submitted gonorrhoea data for the fourth quarter (October to December) 2008.

Of all test-positive data submitted 1.1% were of unknown age and 0.2% were of unknown sex.

Chlamydia Results - October to December, 2008

Chlamydia Incidence

Auckland Region (see table 1)

- Laboratories in the Auckland region tested 36 536 specimens for chlamydia, of which 2 560 (7.0%) specimens tested positive from 2 433 patients.
- Sixty-six percent of all positive patients were aged 15 to 24 years.
- The highest rates of chlamydia in both females and males were found in the 20 to 24 years age group.

Waikato region (see table 1)

- Laboratories in the Waikato region tested 6 885 specimens for chlamydia, of which 715 (10.4%) specimens tested positive from 715 patients.
- Seventy-six percent of all positive patients were aged 15 to 24 years.
- The highest rate of chlamydia was found in the 15 to 19 years age group for females and the 20 to 24 years age group for males.

Bay of Plenty Region (see table 1)

- Laboratories in the BOP region tested 7 161 specimens for chlamydia, of which 743 (10.4%) specimens tested positive from 742 patients.
- Seventy-five percent of all positive patients were aged 15 to 24 years.
- The highest rate of chlamydia was found in the 15 to 19 years age group for females and the 20 to 24 years age group for males.

New laboratories (see table 1)

- New laboratories reported 2 343 (10.3%) test-positive specimens for chlamydia from 2 238 patients.
- Seventy-five percent of all positive patients were aged 15 to 24 years.
- The highest number of chlamydia cases was found in the 15 to 19 years age group for females and 20 to 24 years age group for males.

Data tables: Chlamydia Incidence, October to December, 2008

Table 1.Number of test-positive cases and rates of chlamydia, age group and sex:
October to December 2008

Region	Age group	-					Rate per 100 000			
	(Yrs)	Female	Male	Unknown	Total	Female	Male	Total		
Auckland	<1	18	20	0	38	185.4	196.1	191.0		
	1-9	0	0	0	0	0.0	0.0	0.0		
	10-14	26	5	1	32	52.9	9.6	31.7		
	15-19	607	146	1	754	1223.8	287.5	751.1		
	20-24	656	186	0	842	1298.4	378.3	844.6		
	25-29	257	123	0	380	544.6	284.0	419.9		
	30-34	103	59	0	162	195.5	125.6	162.5		
	35-39	48	39	0	87	85.6	77.5	81.7		
	40+	76	62	0	138	27.4	24.6	26.0		
	Unknown	0	0	0	0	-	-	-		
	Total	1791	640	2	2433	264.7	99.6	184.4		
Waikato	<1	8	3	0	11	324.8	118.8	220.5		
	1-9	0	0	0	0	0.0	0.0	0.0		
	10-14	7	2	0	9	52.9	14.1	32.8		
	15-19	255	61	1	317	1975.8	463.8	1216.4		
	20-24	153	73	1	227	1360.4	628.4	992.9		
	25-29	57	27	0	84	569.9	279.5	427.2		
	30-34	28	14	0	42	247.6	137.2	195.2		
	35-39	6	8	0	14	47.3	70.6	58.3		
	40+	4	7	0	11	5.2	9.9	7.4		
	Unknown	0	0	0	0	-	-	-		
	Total	518	195	2	715	299.3	117.4	210.8		
Bay of Plenty	<1	2	2	0	4	97.6	92.1	94.7		
	1-9	0	0	0	0	0.0	0.0	0.0		
	10-14	23	0	0	23	195.0	0.0	95.6		
	15-19	312	42	1	355	3123.1	398.9	1730.0		
	20-24	157	44	0	201	2156.3	604.8	1380.9		
	25-29	53	24	0	77	681.1	334.4	514.7		
	30-34	21	15	0	36	218.1	177.9	199.4		
	35-39	15	8	0	23	136.0	83.8	111.8		
	40+	15	6		21	20.8	9.2	15.3		
	Unknown	2	0	0	2	-	-	-		
	Total	600	141	1	742	398.5	98.8	253.0		
New	<1	6	5	0	11					
Laboratories	1-9	6	1	1	8					
	10-14	45	2	0	47			/		
	15-19	794	184	0	978		/	/		
	20-24	485	211	1	697					
	25-29	120	101	0	221					
	30-34	57	36	0	93					
	35-39	34	20	0	54					
	40+	25	33	1	59					
	Unknown	33	36		70					
	Total	1605	629	4	2238	/				


Chlamydia trends: Auckland, Waikato and BOP Regions


In the last quarter of 2008, a total of 3 890 cases of chlamydia were reported. This represents a rate of 199.3 per 100 000 population (Auckland: 184.4 per 100 000; Waikato: 210.8 per 100 000; BOP: 253.0 per 100 000).

Compared to the previous quarter, the number of chlamydia diagnoses decreased by 1.1% in the Auckland region, 10.6% in the Waikato region and 0.9% in the BOP region. Chlamydia rates by quarter since January 2003 are shown in Figure 1.

Chlamydia cases reported in this quarter increased by 18.0% on the total of 3 297 in the equivalent quarter of 2007.

The rate of chlamydia in the BOP region was significantly higher than the Waikato and Auckland regions. The rate of chlamydia in the Auckland region was significantly lower than the Waikato region.





Gonorrhoea Results – October to December, 2008

Gonorrhoea Incidence

Auckland Region (see table 2)

- Laboratories in the Auckland region tested 53 119 specimens for gonorrhoea, of which 470 (0.9%) specimens tested positive from 232 patients.
- Sixty percent of all positive patients were aged 15 to 24 years.
- The highest rate of gonorrhoea was found in the 15 to 19 years age group for females and the 20 to 24 years age group for males.

Waikato region (see table 2)

- Laboratories in the Waikato region tested 5 695 specimens for gonorrhoea, of which 47 (0.8%) specimens tested positive from 47 patients.
- Eighty-one percent of all positive patients were aged 15 to 24 years.
- The highest rates of gonorrhoea in both females and males were found in the 20 to 24 years age group.

Bay of Plenty Region (see table 2)

- Laboratories in the BOP region tested 4 718 specimens for gonorrhoea, of which 62 (1.3%) specimens tested positive from 62 patients.
- Seventy-four percent of all positive patients were aged 15 to 24 years.
- The highest rates of gonorrhoea in both females and males were found in the 20 to 24 years age group.

New laboratories (see table 2)

- New laboratories reported 365 (1.2%) test-positive specimens for gonorrhoea from 334 patients.
- Sixty-four percent of all positive patients were aged 15 to 24 years.
- The highest number of gonorrhoea cases was found in the 15 to 19 years age group for females and the 20 to 24 years age group for males.

Data tables: Gonorrhoea Incidence, October to December, 2008

Table 2.Number of test-positive cases and rates of gonorrhoea, age group and sex:
October to December 2008

Region	Age group	Number of test-positive cases				Rate per 100 000		
	(Yrs)	Female	Male	Unknown	Total	Female	Male	Total
Auckland	<1	2	0	0	2	31.1	0.0	15.2
	1-9	0	0	0	0	0.0	0.0	0.0
	10-14	2	1	0	3	6.4	3.0	4.7
	15-19	29	37	0	66	90.9	114.3	102.7
	20-24	25	47	2	74	71.6	141.7	108.7
	25-29	12	24	0	36	37.0	81.2	58.1
	30-34	9	13	0	22	26.1	42.2	33.7
	35-39	2	8	0	10	5.7	25.3	15.0
	40+	3	15	0	18	1.8	9.7	5.6
	Unknown	1	0	0	1	-	-	-
	Total	85	145	2	232	19.8	35.5	27.7
Waikato	<1	0	0	0	0	0.0	0.0	0.0
	1-9	0	0	0	0	0.0	0.0	0.0
	10-14	0	0	0	0	0.0	0.0	0.0
	15-19	8	9	0	17	62.0	68.4	65.2
	20-24	7	14	0	21	62.2	120.5	91.9
	25-29	1	0	0	1	10.0	0.0	5.1
	30-34	1	0	0	1	8.8	0.0	4.6
	35-39	0	3	0	3	0.0	26.5	12.5
	40+	0	4	0	4	0.0	5.7	2.7
	Unknown	0	0	0	0	-	-	-
	Total	17	30	0	47	9.8	18.1	13.9
				T				
Bay of Plenty	<1	0	0	0	0	0.0	0.0	0.0
	1-9	0	0	0	0	0.0	0.0	0.0
	10-14	1	0	0	1	8.5	0.0	4.2
	15-19	6	14	0	20	60.1	133.0	97.5
	20-24	11	15	0	26	151.1	206.2	178.6
	25-29	3	3	0	6	38.6	41.8	40.1
	30-34	2	3	0	5	20.8	35.6	27.7
	35-39	0	1	0	1	0.0	10.5	4.9
	40+	2	0		2	2.8	0.0	1.5
	Unknown	1	0	0	l	-	-	-
	Total	26	36	0	62	17.3	25.2	21.1
New	<1	0	0	0	0			
Laboratories	1-9	2	0	0	2			
	10-14	6	0	0	6			
	15-19	66	46	0	112		/	
	20-24	46	53	2	101			
	25-29	21	31	0	52			
	30-34	6	17	0	23			
	35-39	2	10	0	12	/	/	
	40+	6	16	0	22			
	Unknown	1	1	2	4			
	Total	156	174	4	334			

Gonorrhoea trends: Auckland, Waikato and BOP Regions


In the last quarter of 2008, a total of 341 cases of gonorrhoea were reported. This represents a rate of 23.2 per 100 000 population (Auckland: 27.7 per 100 000; Waikato: 13.9 per 100 000; BOP: 21.1 per 100 000).

Compared to the previous quarter, the number of gonorrhoea diagnoses decreased by 5.7% in the Auckland region, 23.0% in Waikato region and 13.9% in the BOP region.. Gonorrhoea rates by quarter since January 2003 are shown in Figure 2.

Gonorrhoea cases reported in this quarter decreased by 17.2% on the total of 412 in the equivalent quarter of 2007. The largest decrease (20.3%) was seen in the Waikato region (59 in Quarter 4 2007, compared to 47 in Quarter 4 2008).

The rate of gonorrhoea in the Waikato region was significantly lower than the Auckland and BOP regions. However there was no significant difference in the rate of gonorrhoea in the Auckland region compared to the BOP region

Figure 2. Rate of gonorrhoea reported through laboratory surveillance in the Auckland, Waikato and BOP regions: January 2003 to December 2008



Appendix A

STI surveillance data is received from the following laboratories:

- Kaitaia Hospital Laboratory, Northland (Chlamydia only)
- Bay of Islands Hospital Laboratory, Northland (Chlamydia only)
- Northland Pathology Laboratory, Northland
- Whangarei Hospital Laboratory, Northland (Chlamydia only)
- Dargaville Hospital Laboratory, Northland (Chlamydia only)
- North Shore Hospital Laboratory, Waitemata (Chlamydia only)
- LabPlus, Auckland
- Diagnostic Medlab, Auckland
- Middlemore Hospital Laboratory, Counties-Manukau
- Waikato Hospital Laboratory, Waikato
- Medlab Hamilton, Waikato
- Pathlab Waikato, Waikato
- Thames Hospital, Waikato (Chlamydia only)
- Tokoroa Hospital, Waikato (Chlamydia only)
- Te Kuiti Hospital, Waikato
- Taumarunui Hospital, Waikato (Chlamydia only)
- Pathlab Bay of Plenty, Bay of Plenty
- Whakatane Hospital Laboratory, Bay of Plenty
- Rotorua Diagnostic Laboratory, Lakes
- Rotorua Hospital Laboratory, Lakes
- Taupo Southern Community Laboratory, Lakes
- Gisborne Hospital Laboratory, Tairawhiti (Gonorrhoea only)
- Hawke's Bay Hospital, Hawke's Bay
- Hawke's Bay Southern Community Laboratory, Hawke's Bay
- Taranaki MedLab, Taranaki
- Medlab Central, MidCentral
- Hutt Hospital Laboratory, Hutt Valley
- Aotea Pathology, Capital & Coast (Gonorrhoea only)
- Grey Hospital Laboratory, West Coast
- Canterbury Health Laboratories, Canterbury (Gonorrhoea only)
- Christchurch Southern Community Laboratory, Canterbury
- Ashburton Southern Community Laboratory, Canterbury
- Oamaru Southern Community Laboratory, Otago
- Otago Southern Community Laboratory, Otago
- Balclutha Southern Community Laboratory, Otago
- Dunstan Southern Community Laboratory, Otago
- Queenstown Southern Community Laboratory, Southland
- Gore Southern Community Laboratories, Southland
- Invercargill Southern Community Laboratory, Southland