

National Wastewater Surveillance Programme - COVID-19

Week 32 (Week Ending 14 August 2022)

Report prepared on 17 August 2022

96%

Of sites tested had SARS-CoV-2 detected (95/99 sites).

73%


Of the NZ population covered by wastewater testing

Omicron BA.4/5 (~98%)

Most prevalent variant detected BA.2 (~2%)

Overall, SARS-CoV-2 levels in wastewater have decreased

Omicron BA.4/5 levels continue to increase relative to BA.2



All data reported as a 7-day average

Figure 1. National timeseries of estimated SARS-CoV-2 genome copies (GC) in wastewater rate (GC/person/day) and reported case rate (per 100,000 population per day). Numbers in the points are the week of the year. The most recently reported weeks are highlighted in green. Log10 scale.

Results For Week 32 2022

In the week ending 14 August 2022, 147 samples were collected from 99 locations in New Zealand.

SARS CoV-2 RNA was **detected** in 142 samples from 95 (97%) sites tested this week (Figure 2, Figure 3 and Table 1). SARS-CoV-2 was **not detected** in samples collected from Eketāhuna (collected 9/8), Kaponga (9/8), Porangahau (10/8), Reefton (8/8) and Waipukurau (not detected on 8/8 but detected on 11/8). A total of 7 samples were below the limit of quantification.






Figure 2. Results for SARS-CoV-2 RNA in wastewater collected across New Zealand.

For the week ending **14 August 2022**, **26**% of sites have **increased** SARS-CoV-2 levels compared to the previous week and **47**% have **decreased** levels. Compared to a month ago, **20**% of sites show an **increase** and **63**% of sites show a **decrease** (Figure 3).



Figure 3. Comparison of SARS-CoV-2 levels for the week ending 14 August 2022, compared with the levels measured: **A) 1 week ago**; **B) 2 weeks ago**; **C) 4 weeks ago**; **D) 12 weeks ago**. Only sites with results for both time points are included. When the viral quantity is 30% or more higher this is labelled as increased (red up arrow on map). When the viral quantity is 30% or more lower this is labelled as decreased (blue down arrow on map). If viral levels have changed less than this in the compared weeks, this is labelled as no change (yellow circle on map). Interactive map of weekly results available publicly at https://www.esr.cri.nz/our-expertise/covid-19-response/wastewater-testing-results

Regional Trends

Regional summaries (Figure 4) also indicate regional decreases in genome copies per person per day (rolling 2-week simple average) across all regions except for the Northern region compared with the previous week.




Figure 4 The average SARS-CoV-2 genome copies detected per person per day in the five Ministry of Health regions. Weighted by population tested that week in each region.

Wastewater Variant Analysis

Consistent with whole genome sequencing (WGS) of clinical cases, the BA.4 and BA.5 variants are the dominant circulating variants across Aotearoa (national average of ~98%, Figure 5). This week, 13/20 sites were 100% BA.4/5 (i.e BA.2* could not be detected).

The displacement of BA.2 has been rapid - over the past 10 weeks nationally (aggregated across all sentinel sites,) BA.4/5 was 2.8%, 6%, 10.5%, 33%, 53%, 73%, 81%, 90%, 93% **and now 98% (**Fig 6**).** There were no detections of BA.1 this week.




Figure 5. Variants present in New Zealand wastewater from 20 sentinel sites during week 31 (8-14 August 2022).






Figure 6. Change in variant prevalence over time at a national scale. Data is collected from up to 21 sentinel sites each week.


Interpreting site graphs





- 1 Site Name
- Sample Collection method and population. Results based on autosampler may be more representative than grab sample-based results.
- Wastewater Results Shown as Solid Line | Rolling 14-day average of genome copies/person/day on a log10 scale.
- 4 Individual Results Samples Shown as Circles | Rolling 14-day average of genome copies/person/day on a log10 scale.
- Rolling 7-day average new cases shown as dashed line | New cases reported in a catchment based on reported date of illness on a log10 scale. This data is not available for all sites & subject to change.


Note: Wastewater and cases data are on a log10 scale. Scales on all graphs have been normalised to cover the same scale on every graph. Care should be taken when interpreting the data.












































10K

1K


100

10


10K

10

Jul










Table 2. Weekly summary of results

Key: Not Detected Detected (below limit of quantification) Detected) Not sampled

· '	2022	2																										
Site		06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32
Northland					_		_	-	-		-					-	-		-	_	-		_		-	-	-	
Ahipara																												
Dargaville																												
Haruru																												
Hikurangi																												
Kaeo																												
Kaikohe																												
Kaitaia																												
Kaiwaka																												
Kawakawa																												
Kerikeri																												
Kohukohu																												
Mangawhai																												
Maungaturoto																												
Opononi																												
Paihia																												
Rawene																												
Ruakaka																												
Russell																												
Taipa																												
Whangarei																												
Whatuwhiwhi																												
Auckland				'			_		'	'										'		'				_		
Army Bay																												
Auckland																												
(Combined)																												
Auckland East																												
Auckland Southwest																												
Auckland West																												
Beachlands																												
Clarks Beach																												

	2022	2																										
Site		06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32
Helensville																												
Kawakawa Bay																												
Kingseat																												
North Shore																												
Omaha																												
Pukekohe																												
Snells/Algies																												
Waiuku																												
Warkworth																												
Wellsford																												
Waikato					•	•		'				•	•		•	·	'	•	•		•		•	,	•		•	
Cambridge																												
Hamilton																												
Matamata																												
Morrinsville																												
Ngatea																												
Otorohanga																												
Paeroa																												
Putaruru																												
Taupo																												
Te Awamutu																												
Thames																												
Tokoroa																												
Turangi																												
Waihi																												
Whangamata																												
Whitianga																												
Bay of Plenty					'																							
Katikati																												
Kawerau																												
Maketu																												
Mt Maunganui																												
Opotiki																												
Rotorua																												
Tauranga																												

	2022	2																										
Site		06	07	80	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32
Te Puke																												
Waihi Beach																												
Whakatane																												
Gisborne					'	'	'			'	,	•	•		'	·	'	•		'		•		·	'	•		•
Gisborne																												
Hawke's Bay			'		'		'		'		'	'	'	'	'	'		'	'		'		'	'		'		
Hastings																												
Mahia																												
Napier																												
Otane																												
Porangahau																												
Takapau																												
Te Paerahi																												
Waipawa																												
Waipukurau																												
Wairoa																												
Taranaki						'	'			'	,	•	•		'	·	'			'				·	'	•		
Eltham																												
Hawera																												
Kaponga																												
Manaia																												
New Plymouth																												
Opunake																												
Patea																												
Stratford																												
Waverley																												
Manawatu-																												
Whanganui																												
Dannevirke																												
Eketahuna																												
Feilding																												
Levin																												
Pahiatua																												
Palmerston North																												

Site 05 06 07 08 09 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 3.		2022	2																										
What woodville Wellington Carterton Featherston Greytown Hutt Valley Masterton Otaki Paraparaumu Porirua Wellington (Moa Polinty) Wellington We	Site			07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32
Urban Wellington Carterton Featherston Greytown Hutt Valley Martinborough Masterton Greytown Hutt Valley Masterton Greytown Hutt Valley Masterton Greytown Hutt Valley	Taumarunui																												
Woodwile Wellington	Whanganui																												
Wellington Carterton Featherston Greytown Hutt Valley Martinborough Masterton Otaki Paraparaumu Porirua Wellington (Moa Point) Wellington (Western) Tasman Motueka Nelson Nelson Central/North Richmond/Nelso n South Martiborough Blenheim Picton West Coast Greymouth																													
Carterton Featherston Greytown Hutt Valley Martinborough Masterton Otaki Paraparaumu Porirua Wellington (Moa Point) Wellington (Western) Tasman Motueka Nelson Nelson Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth																													
Featherston Greytown Hutt Valley Martinborough Masterton Otaki Paraparaumu Porirua Wellington (Moa Point) Wellington (Western) Tasman Motueka Nelson Nelson Nelson Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth	Wellington																												
Greytown Hutt Valley Martinborough Masterton Otaki Paraparaumu Porirua Wellington (Moa Polint) Wellington (Moa Polint) Wellington (Western) Tasman Motueka Nelson Nelson Central/North Richmond/Nelso n South Mariborough Blenheim Picton West Coast Greymouth	Carterton																												
Hutt Valley Martinborough Masterton Otaki Paraparaumu Porirua Wellington (Moa Point) Wellington (Western) Tasman Motueka Nelson Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth	Featherston																												
Masterton Otaki Paraparaumu Porirua Wellington (Moa Point) Wellington (Western) Tasman Motueka Nelson Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth	Greytown																												
Masterton Otaki Paraparaumu Porirua Wellington (Moa Point) Wellington (Western) Tasman Motueka Nelson Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth	Hutt Valley																												
Otaki Paraparaumu Porirua Wellington (Moa Point) Wellington (Western) Tasman Motueka Nelson Nelson Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth	Martinborough																												
Paraparaumu Porirua Wellington (Moa Point) Wellington (Western) Tasman Motueka Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth	Masterton																												
Porirua Wellington (Moa Point) Wellington (Western) Tasman Motueka Nelson Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth	Otaki																												
Porirua Wellington (Moa Point) Wellington (Western) Tasman Motueka Nelson Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth	Paraparaumu																												
Point) Wellington (Western) Tasman Motueka Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth																													
Point) Wellington (Western) Tasman Motueka Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth	Wellington (Moa																												
(Western) Tasman Motueka Nelson Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth	Point)																												
Tasman Motueka Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth	Wellington																												
Motueka Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth																													
Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth																													
Nelson Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth																													
Central/North Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth																													
Richmond/Nelso n South Marlborough Blenheim Picton West Coast Greymouth																													
n South Marlborough Blenheim Picton West Coast Greymouth																													
Blenheim Picton West Coast Greymouth	n South																												
Picton West Coast Greymouth	Marlborough																												
West Coast Greymouth	Blenheim																												
Greymouth	Picton																												
	West Coast																												
	Greymouth																												
	Hokitika																												
Reefton	Reefton																												
Westport																													
Canterbury																								1					
Amberley																													
Ashburton																													
Christchurch																													

	2022	2																										
Site	05	06	07	80	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32
Hanmer Springs																												
Kaiapoi																												
Kaikoura																												
Leeston																												
Oamaru																												
Rangiora																												
Rolleston &																												
Eastern Selwyn																												
Timaru																												
Woodend																												
Otago																												
Alexandra																												
Balclutha																												
Cromwell																												
Dunedin (Green																												
Island)																												

Overview

SARS-CoV-2, the virus that causes COVID-19 disease, is shed in the faeces of people that are infected and so the viral RNA can be detected in wastewater. As such, testing wastewater for SARS-CoV-2 RNA is an efficient population-based COVID-19 surveillance tool. Based on national and international data, this method has been shown to be an indicator of increasing and decreasing cases (i.e., early warning system) and complements other surveillance tools. A national wastewater COVID-19 surveillance programme was established in 2021 by the Institute of Environmental Science and Research (ESR). This work is funded by the New Zealand Ministry of Health and is part of New Zealand's COVID-19 response.

Wastewater samples are collected from wastewater treatment plants across both the North and South Island of New Zealand. Grab or 24hr composite samples are collected. Most sites are sampled at least weekly between Monday and Thursday of any given week. The number of sites and frequency of collection varies over time. Greater variability is expected with grab samples.

Approach

Samples are sent from each wastewater treatment plant to one of the ESR laboratories (Porirua or Christchurch). Processing involves the concentration of virus and extraction of viral RNA. The presence of SARS-CoV-2 RNA in the sample is then determined using RT-qPCR.

A result of **not detected**, means that SARS-CoV-2 RNA is either absent from the sample, or at a level too low to be detected. When SARS-CoV-2 RNA is **detected**, the concentration in the sample can be calculated. Low amounts of SARS-CoV-2 RNA in a sample may not be able to be accurately quantified and are recorded as less than the limit of quantitation. For quantitation, the raw concentration data (i.e., genome copies per L) is converted to a viral load of **genome copies per day per person**. This calculation considers the flow rate of wastewater entering the wastewater treatment plant and the population in the catchment. This is the population-normalised viral load.

Key Points & Limitations

- SARS-CoV-2 RNA concentrations should not be compared between wastewater catchments.
- Day to day variability in SARS-CoV-2 RNA concentrations, especially in smaller catchments, is to be expected. Greater variability is expected with grab samples.
- Generally, increasing viral loads are associated with increasing numbers of people with SARS-CoV-2 infection and vice versa (decreasing concentrations indicating decreasing cases). However, there are many factors that affect the amount of viral RNA detected and so data from wastewater surveillance cannot indicate the exact number of COVID-19 cases in the catchment area.
- The number of COVID-19 cases reported via individual testing are reported for each region to
 provide a comparison to the wastewater results. The cases in each catchment area are an estimate
 of the number of people in that wastewater catchment area that have reported a positive test.
 However, because the wastewater catchments do not exactly align with regional boundaries, the
 number of cases estimated by region and by water catchment area may be different.
- Data are provisional and may be subject to change by location.
- As septic tank systems are not connected to wastewater treatment plants, the wastewater from these households will not be represented in the data.

Acknowledgements

This work represents the combined efforts of many individuals and organisations.

We continue to be indebted to the teams across the country who are collecting the wastewater that underpins this work.

The wastewater analysis has been undertaken at the ESR Kenepuru (in Porirua) and Christchurch laboratories by a team which may on any given week include contributions from: Kirsten Carter, Joanne Chapman, Dawn Croucher, Joanne Hewitt, Joycelyn Ho, Anower Jabed, Susan Lin, Olivia Macrae, Ashley McDonald, Andrew Ng, Ashley Orton, Paula Scholes and Fatiha Sulthana. Data science analysis, visualisation and reporting is the result of team effort from: Franco Andrews, Bridget Armstrong, Raewyn Campbell, Gerhard de Beer, Richard Dean, Brent Gilpin, Joanne Hewitt, Dawen Li, Helen Morris and Bindu Priya. Ongoing support for this work from the Ministry of Health and ESR management is appreciated.

Notes

Sites and frequency of sample collection: The catchment population sites selected for the surveillance range from approximately 100 to over 1,000,000 individuals. The sites cover all regions of the country. Most major towns and all cities, as well as many smaller communities, are included. In early 2022, the wastewater catchment areas cover over 80% of the population connected to wastewater treatment plants. The sites from which samples have been collected have varied over the last 12 months. New sites may be added over time, and/or sampling may reduce in frequency or cease for other sites. The selection and frequency of sampling vary depending on the local population, access to wastewater collection points, staff availability to collect samples and risk factors. When included, samples are collected at least weekly, with twice weekly sampling being common. A number of samples have also been collected from non-WWTP sites (manholes and pump stations- mostly in Auckland).

Sampling method: The preferred option is to automatically collect a 24 hour 'composite' sample. This is where a pump automatically collects a small volume of wastewater every 15 minutes over 24 hours using a composite sampler. These samplers are available in some wastewater treatment plants. When composite samplers are not available, 'grab' samples are collected. These range from a sample being taken at a single point in time, to 3 samples taken over 30 minutes, to samples collected over a day. Grab samples represent only the composition of the source at that time of collection and may not be as representative as a 24-hour composite sampler. More variation may be expected with grab samples.

Laboratory analysis of wastewater samples: Samples are sent from each wastewater treatment plant to one of the ESR laboratories (Porirua or Christchurch). Processing of each sample commences within an hour or two of receipt. Processing involves the concentration of virus from 250 mL sample to approx. 1 mL using centrifugation and polyethylene glycol. Viral RNA is then extracted from a small volume of 0.2 mL concentrate to give a final volume of 0.05 mL The presence of SARS-CoV-2 RNA is determined using RT-qPCR. SARS-CoV-2 is considered detected when any of the RT-qPCR replicates are positive.

RT-qPCR: Reverse transcription (RT) to convert RNA to complementary DNA (cDNA), followed by quantitative PCR (qPCR). RT-qPCR is used for detection and quantification of viral RNA.

Method sensitivity: The protocol used to concentrate SARS-CoV-2 from wastewater allows for the sensitive detection of SARS-CoV-2 by RT-qPCR. ESR has shown that when 10 individuals are actively shedding SARS-CoV-2 RNA in a catchment of 100,000 individuals, there was a high likelihood of detecting viral RNA in wastewater (https://doi.org/10.1016/j.watres.2021.118032). Shedding by one individual may be detected in wastewater, but it does depend on many factors including the amount and duration of shedding. Very low levels in wastewater may be not able to be quantified (i.e., less than the limit of quantification- see below).

SARS-CoV-2 RNA detected (positive result): A positive detection in the wastewater indicates that at least one person has been shedding SARS-CoV-2 into the wastewater at some point during the time period that the sample was being collected. In some cases, detections could also be due to the shedding of low levels of SARS-CoV-2 RNA by a recently recovered case. The detection of SARS-CoV-2 RNA does not indicate that infectious virus is present.

SARS-CoV-2 RNA not detected (negative result): A negative result can occur because there are no active 'shedding' cases in the catchment or because the SARS-CoV-2 RNA concentration is too low to be detected, most likely because there are a very low number of cases in the wastewater catchment. Therefore, negative finding does not necessarily guarantee the absence of COVID-19 in the community.

Viral loads and normalisation: When detected, the SARS-CoV-2 RNA concentration is calculated as genome copies per L of wastewater. This is then converted to a viral load of **genome copies/day/person**. This conversion takes into account the flow rate of wastewater entering the treatment plant (the influent) and the population in the catchment. The **flow rate** is the total volume (m3 per day) recorded at the inlet of the wastewater treatment plant over 24 hours. This is a **population-normalised viral load**. Currently, the flow rate is the average annual flow rate, but will be replaced with daily flow rate when available (note that rainfall may significantly increase the flow rate at the inlet, diluting the sample, and may result in lower concentrations and a false negative result).

In future, SARS-CoV-2 RNA concentrations will also be normalised by testing for the presence of pepper mild mottled virus (PMMoV). PMMoV is a virus that infects peppers but not humans. Consumption of peppers or pepper products, such as chilli sauce, means that PMMoV is detected in wastewater – normally at very high concentrations. Therefore, PMMoV has been found to be a useful proxy for the amount of faecal material in a wastewater sample. For normalisation, the concentration of SARS-CoV-2 RNA is divided by that of PMMoV in each sample. Different normalisation methods may result in changes to some data points, but trends are unlikely to change significantly.

Limit of quantification: The lowest concentration of the target that can be reliably quantified is referred to as the limit of quantification. For those samples where SARS-CoV-2 is detected but cannot be quantified, a value of 5 genome copies/mL wastewater is used. While a standard method is being used, virus recovery can vary from sample to sample, and this may affect the quantitation.

Data subject to change: Data generated for the New Zealand Wastewater COVID-19 Surveillance Programme should be considered provisional and may be subject to change. Data may be incomplete for the most recent 2-week period due to processing, testing and reporting delays.

Data not shown:

- Data from 'ad hoc' sampling locations including from individual facilities/building (e.g., workplaces, prisons, MIQs) are not included.
- Results from certain samples may not be shown, as the result was either deemed invalid, or the sample could not be tested (e.g., leaked in transit, not labelled).

For further information please contact

Brent Gilpin Joanne Hewitt Science Leader Senior Scientist, Environmental & Food Virology

Brent.gilpin@esr.cri.nz Joanne.hewitt@esr.cri.nz